

A LUMINOUS RECURRENT SUPERSOFT X-RAY SOURCE IN NGC 300

A.K.H. KONG¹ AND R. DI STEFANO^{1,2}
Draft version February 2, 2008

ABSTRACT

We report the results of *XMM-Newton* observations for an especially luminous supersoft X-ray source (SSS) with bolometric luminosity of 10^{39} erg s⁻¹ in the spiral galaxy NGC 300. The source was detected as a SSS in 1992 and disappeared in subsequent X-ray observations. The source was active again during recent *XMM-Newton* observations. It appeared to be very soft ($kT \sim 60$ eV) and very luminous ($\sim 10^{38} - 10^{39}$ erg s⁻¹). The two *XMM-Newton* observations also reveal that the source went from a “high” state to a “low” state in 6 days. We also found a 5.4-hr periodicity during the “low” state. We consider white dwarf, black hole, and neutron star models to explain the nature of the source.

Subject headings: galaxies: individual (NGC 300) — X-rays: binaries — X-rays: galaxy

1. INTRODUCTION

Supersoft X-ray sources (SSSs) form a distinct class of objects, first established through *ROSAT* observations. The hallmarks of SSSs are very soft X-ray emission (kT typically < 100 eV) and bolometric luminosities of $10^{36} - 10^{38}$ erg s⁻¹. The advent of *Chandra* and *XMM-Newton* provides good opportunities to detect and study SSSs in nearby galaxies. Luminous (10^{38-40} erg s⁻¹) SSSs have been found in several nearby galaxies including M31 (Kong et al. 2002; Di Stefano et al. 2002a), M81 (Swartz et al. 2002), M101 (Pence et al. 2001; Mukai et al. 2003; Di Stefano & Kong 2003 [DK03]), NGC4697 (Sarazin, Irwin & Bregman 2001; DK03), M51 (DK03), and M83 (Soria & Wu 2003; DK03).

NGC 300 is an SA(s)d galaxy, seen near face-on (inclination angle 46° ; Tully 1988) at a distance of 2.0 ± 0.1 Mpc (Freedman et al. 2001). The galaxy has been observed by *ROSAT* (Read et al. 1997; Read & Pietsch 2001) and *XMM-Newton* (Soria & Kong 2003).

In this Letter, we report the reappearance of a luminous SSS in NGC 300 as observed with *XMM-Newton*.

2. OBSERVATIONS AND DATA REDUCTION

2.1. ROSAT

NGC 300 was observed by *ROSAT* five times from 1991 to 1997. A detailed analysis of the *ROSAT* data was done by Read & Pietsch (2001). Briefly, the datasets consist of 2 Position Sensitive Proportional Counter (PSPC) and 3 High Resolution Imager (HRI) pointings. The exposures range from ~ 9 ks to ~ 37 ks. The luminous supersoft source was only detected in a 37 ks PSPC observation taken on 1992 May and June (see Read & Pietsch 2001). Spectral analysis was also done by Read et al. (1997). We have reanalyzed the PSPC spectrum taken on 1992 May/June, and used other data to set upper limits on the luminosity for the long-term lightcurve.

We extracted the source spectrum from a $30''$ circular region, while background was from an annulus region ($45''$ and $60''$ radii) centered on the source. The spectrum was grouped into at least 20 counts per spectral bin to allow χ^2 statistics to be used.

2.2. XMM-Newton

The *XMM-Newton* instrument modes were full-frame, medium filter for the three European Photon Imaging Cameras (EPIC). The first observation was taken on 2000 December 26 for 37 ks and the second observation was on 2001 January 1 for about 47 ks. After rejecting intervals with a high background level, we considered a good time interval of ~ 28 ks and ~ 40 ks for the first and second observation, respectively. Data were reduced and analyzed with the *XMM-Newton* SAS package v5.3.3.

We used here the MOS images to determine the position of the source because the spatial resolution of MOS detector is slightly better than that of pn. The source is located at R.A.=00h55m10s.7, Dec.= $-37^\circ 38' 55''$ (J2000), $\sim 4'$ (2.4 kpc) from the galactic center; the derived positions from the two *XMM-Newton* observations agree with each other. This position is about $5''$ off from previous *ROSAT* PSPC observation, and is consistent with the positional error ($7.3''$) quoted by Read & Pietsch (2001).

Source spectra and lightcurves of XMMU J005510.7-373855 were extracted with the SAS task XMMSELECT. Source-free regions were used for background to avoid the chip boundary and a nearby faint source. In order to allow χ^2 statistics to be used, all the spectra were binned such that there are at least 20 counts per spectral bin. Response matrices were created by RMFGEN and ARFGEN.

3. ANALYSES AND RESULTS

Spectral analysis was performed by making use of XSPEC v11.2. Table 1 shows the best-fitting spectral parameters for the three *ROSAT* and *XMM-Newton* observations.

For the *ROSAT* observation, the spectrum (see Figure 1) can be fit with a blackbody model with $N_H = 1.1 \times 10^{21}$ cm⁻² and $kT = 48.7$ eV; the 0.2–2 keV luminosity is 10^{39} erg s⁻¹. We note that Read et al. (1997) fit the spectrum with a thermal bremsstrahlung model with temperature of a 0.1 keV and we confirmed that it is also an acceptable model. Using the thermal bremsstrahlung model, the luminosity becomes 3.8×10^{38} erg s⁻¹ [the luminosity quoted by Read et al. (1997) is 1.6×10^{37} erg s⁻¹, which is an absorbed luminosity corrected for absorption in our own Galaxy], a factor of 2.5 lower than the blackbody model. While we cannot distinguish between

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138; akong@cfa.harvard.edu

² Department of Physics and Astronomy, Tufts University, Medford, MA 02155

the two models statistically, we prefer the blackbody model as subsequent *XMM-Newton* observations confirm the super-soft nature of the source (see below). Also, the *XMM-Newton* observations would have detected the high energy photons associated with the thermal bremsstrahlung model, had the model been correct. Finally, the *XMM-Newton* observations provide more photons than *ROSAT*.

We fit the pn, MOS1 and MOS2 data simultaneously with several single-component models with interstellar absorption (including absorbed power-law, thermal bremsstrahlung, blackbody and Raymond-Smith); only the blackbody model provides an acceptable fit. The blackbody temperature of both observations ranges between 57 eV and 67 eV, while the N_H varies from $1.5 \times 10^{21} \text{ cm}^{-2}$ in the first observation to $5.3 \times 10^{20} \text{ cm}^{-2}$ in the second observation. The 0.2–2 keV luminosity also drops from $9 \times 10^{38} \text{ erg s}^{-1}$ to $1.2 \times 10^{38} \text{ erg s}^{-1}$, indicating that the source is a variable on timescale of days. The spectra of the two observations are shown in Figure 1.

The combined *ROSAT* and *XMM-Newton* long-term lightcurve of the SSS is shown in Figure 2, which is constructed from a series of *ROSAT* and *XMM-Newton* pointings. The source was below the detection limit of other *ROSAT* observations. In these cases, we estimated the 3σ limits, assuming a blackbody model with mean N_H (10^{21} cm^{-2}) and kT (58 eV). The value of N_H plays an important role in estimating the luminosity. For instance, if we lower the N_H to $5.3 \times 10^{20} \text{ cm}^{-2}$, as found in the second *XMM-Newton* observation, the upper limits decrease by a factor of ~ 3 . In other words, the source varies by as much as factor of 30 between the “low” state and the “high” state spanning 8.5 years.

To search for short-term variability, we computed the Lomb-Scargle periodogram (LSP; Lomb 1976; Scargle 1982), a modification of the discrete Fourier transform which is generalized to the case of uneven spacing. In each observation, we extracted the combined background subtracted lightcurve from pn and MOSs to increase the signal-to-noise. Individual lightcurves from each of the three detectors were also used to verify the result. By applying the LSP to the combined lightcurve, we found that there is a sharp peak at 5.4 hr in the “low” state observation (see Figure 3). Independent checks from the pn and MOS data also confirmed the periodicity. We determined the 99.9% confidence level by generating Gaussian noise datasets with the same time intervals and variance as the true data, and then performed the LSP analysis on the resulting datasets (see Kong, Charles & Kuulkers 1998). The peak power in each periodogram (which must be purely due to noise) was then recorded. This was repeated 10000 times to obtain good statistics. The peak at 5.4 hr is well above the 99.9% confidence level. The folded lightcurve of the “low” state data in 5.4 hr is also shown in Figure 3.

Similar analysis was also carried out in the “high” state data, but there is no significant peak in the LSP. The folded lightcurve shows no obvious periodic variability at 5.4 hr (see Figure 3). It is not clear if it is due to geometric effect but the exposure time of the “high” state covers less than 8 hr, corresponding to 1.4 cycles of the 5.4-hr period in contrast to the 2.1 cycles in the “low” state. We therefore performed Kolmogorov-Smirnov test to test for variability and found significant variability at the 99.9% levels. We also search for variability of the hardness ratio but no significant change was found in both observations.

We also examined an Digitized Sky Survey image of NGC

300 to search for possible optical counterpart. Within a $4''$ error circle of the *XMM-Newton* position, there is no obvious optical counterpart with limiting B and R magnitudes of 21 and 18.1, respectively. However, the source is surrounded by HII regions (Deharveng et al. 1988; Blair & Long 1997), OB associations (Pietrzynski et al. 2001) and supernova remnants (Blair & Long 1997). The nearest HII region (source 136; Deharveng et al. 1988) is located about $20''$ (~ 200 pc) south-east of the X-ray source. Within $50''$ (~ 500 pc) of the SSS, we found an OB association (AS99; Pietrzynski et al. 2001), two supernova remnant candidates (N300-S21, N300-N23; Blair & Long 1997) and a HII region (N300-H20; Blair & Long 1997). We also looked for UV counterpart from the Optical Monitor (OM) image (see Soria & Kong 2003 for details of the OM observations); no counterpart is found at the position of the SSS in both observations.

4. DISCUSSION

It is clear from the spectral fits of all *ROSAT* and *XMM-Newton* observations that the XMMU J005510.7-373855 is very soft ($kT \sim 48 - 67$ eV) and is highly variable. Such a soft spectrum is consistent with SSSs seen in our own Galaxy, the Magellanic Clouds, M31 (Greiner 2000³) and several nearby galaxies (see § 1). Since the Galactic and Magellanic Cloud SSSs are conjectured to be white dwarfs (WDs), it is reasonable to consider a WD model for the source in NGC 300. Since, however, the luminosity in the high state may be too large to be consistent with the standard SSS WD model, we also consider black hole (BH) and neutron star (NS) models.

The X-ray emission of the SSS could come from a WD that burns accreting hydrogen in a quasi-steady manner. The maximum luminosity is the Eddington luminosity for a $1.4M_\odot$ object. WD models therefore seem to be ruled out by the high bolometric luminosities during 2 of the observations. It is nevertheless worth noting that the WD model temperatures are comparable to those we observed, and that X-ray variability by a factor of a few over times of days and months has been observed in SSSs. (see e.g., Greiner and Di Stefano 2002 and references therein.) If the 5.4-hour variation detected during the X-ray low state corresponds to an orbital period, then this system may be similar to 1E0035.4-7230, a Magellanic Cloud SSS that appears to have a 4 hour period (Kahabka 1996). In such a short-period binary, it is likely that a radiation-driven wind plays an important role in the binary evolution. (see, e.g., van Teeseling & King 1998) A WD model would perhaps be the most natural explanation for the SSS in NGC 300, were it not for the high estimated bolometric luminosities. Note that it may be difficult to use beaming to circumvent the Eddington limit in these systems, since the inner region of the accretion disk can also become a luminous emitter of SSS radiation (Popham & Di Stefano 1996).

SSS emission is expected from accreting BHs. Modeling the accretion disk as a thin disk which is optically thick, we can derive a relationship between the minimum mass of the accretor, and the observed temperature and luminosity (Frank, King, & Raine 2002). Using $kT = 60$ eV and $L_X = 10^{38} \text{ erg s}^{-1}$ ($10^{39} \text{ erg s}^{-1}$), we find that the accretor mass is greater than $\approx 890M_\odot$ ($2800M_\odot$). Although the fact that the observed luminosity is $10^2 - 10^3$ times smaller than the Eddington luminosity may call the validity of the disk model into question, this calculation suggests that the accretor is an intermediate-mass

³ see <http://www.aip.de/~jcg/sss/ssscat.html> for updated catalog

BH. If the accretor is a BH, we may use $L_{\text{obs}} = 0.1\dot{m}c^2(\eta/0.1)$, where $\dot{m}$ is the accretion rate and η , the efficiency factor, is likely to be close to 0.1. This yields $\dot{m} \approx 1.8 \times 10^{-8} M_{\odot}/\text{yr}$ ($1.8 \times 10^{-7} M_{\odot}/\text{yr}$) for $L_{\text{obs}} = 10^{38} \text{ erg s}^{-1}$ ($L_{\text{obs}} = 10^{39} \text{ erg s}^{-1}$). This would be consistent with an irradiation-driven wind from a low-mass donor. Measurement of the orbital period could provide supporting evidence. Indeed, if the donor fills its Roche lobe, then $P_{\text{orb}} \approx (8.9 \text{ hrs})(M_d/M_{\odot})$, where M_d is the mass of the donor star. If $P_{\text{orb}} = 5.4 \text{ hrs}$, then $M_d \approx 0.61 M_{\odot}$.

It is of course possible that the observed emission emanates from an accreting NS. In this case the SSS emission would presumably emanate from a photosphere which is much larger than the neutron star itself. The photospheric radius would be different in each observation: $3.7 \times 10^9 \text{ cm}$ during the *ROSAT* observation, $2.6 \times 10^9 \text{ cm}$ during the *XMM-Newton* high-state observation, and $6.8 \times 10^8 \text{ cm}$ during the *XMM-Newton* low-state observation, corresponding to the different temperatures and luminosities.

At this point, neutron star models seem conceptually unattractive because there is no obvious explanation for why the photosphere should achieve these relatively large sizes, and no way to relate them to the system's fundamental physical parameters. In addition, just as in the WD models, the luminosity appears to be super-Eddington during two of the observations, unless the neutron star has a mass as large as $3-4 M_{\odot}$. In this case, if the efficiency of turning accretion energy into X-rays is the same during all 3 observations, then the accretion

rate must have changed by an order of magnitude, approaching $2 \times 10^{-8} M_{\odot} \text{ yr}^{-1}$ during the high state. (see Di Stefano et al. 2002b for a similar model of the X-ray source in the globular cluster Bo 375.) If, on the other hand, the mass of the neutron star is close to $1.4 M_{\odot}$, then other ways around the Eddington limit must be found; e.g., the energy could be beamed into a smaller solid angle during the 2 high-state observations. In this case, the accretion rate could be significantly lower.

5. SUMMARY

We have studied a recurrent luminous SSS in NGC300 with *XMM-Newton*. The source was seen by *ROSAT* in 1992 and fell below the detection limit in subsequent *ROSAT* observations. It reappeared in recent *XMM-Newton* observations with bolometric luminosities between 10^{38} and $10^{39} \text{ erg s}^{-1}$. During the “low” state, the source showed a 5.4-hr periodicity. If we consider the periodicity is due to the orbital period, then the X-ray emission can be explained by WD, BH, and NS models. However, WD and NS models appear to be unlikely due to the high X-ray luminosity during the “high” state. Further repeated X-ray observations of the SSS in different states may permit discrimination among these models.

This work was supported by NASA under an LTSA grant, NAG5-10705. A.K.H.K. acknowledges support from the Croucher Foundation.

REFERENCES

- Blair, W.P., & Long, K.S. 1997, *ApJS*, 108, 261
 Deharveng, L., Caplan, J., Lequeux, J., et al. 1988, *A&AS*, 73, 407
 Dickey, J.M., & Lockman, F.J. 1990, *ARA&A*, 28, 215
 Di Stefano, R., & Kong, A.K.H. 2003, *ApJ*, in press (astro-ph/0301162), DK03
 Di Stefano, R., Greiner, J., Kong, A., Garcia, M.R., Primini, F.A., Barmby, P., Murray, S.S., Curry, S. 2002a, *American Physical Society*, APRN17072
 Di Stefano, R., Kong, A.K.H., Garcia, M.R., Barmby, P., Greiner, J., Murray, S.S., & Primini, F.A. 2002b, *ApJ*, 570, 618
 Frank, J., King, A., & Raine, D.J. 2002, *Accretion Power in Astrophysics: Third Edition*, CUP
 Freedman, W.L., et al. 2001, *ApJ*, 553, 47
 Greiner, J. 2000, *NewA*, 5, 137
 Greiner, J., & Di Stefano, R. 2002, *ApJ*, 578, L59
 Kahabka, P. 1996, *A&A*, 306, 795
 Kong, A.K.H., Charles, P.A., & Kuulkers, E. 1998, *NewA*, 3, 301
 Kong, A.K.H., Garcia, M.R., Primini, F.A., Murray, S.S., Di Stefano, R., & McClintock, J. 2002, *ApJ*, 577, 738
 Lomb, N.R. 1976, *Ap&SS*, 39, 447
 Mukai, K., Pence, W.D., Snowden, S.L., Kuntz, K.D. 2003, *ApJ*, 582, 184
 Pence, W.D., Snowden, S.L., Mukai, K., & Kuntz, K.D. 2001, *ApJ*, 561, 189
 Pietrzynski, G., Gieren, W., Fouque, P., & Pont, F. 2001, *A&A*, 371, 497
 Popham, R., & Di Stefano, R. 1996, in *Supersoft X-Ray Sources*, Proceedings of the International Workshop Held in Garching, Germany, 28 February - March 1996. Lecture Notes in Physics, Vol. 472, edited by Jochen Greiner. Springer-Verlag, Berlin Heidelberg New York, p.65
 Read, A.M., Ponman, T.J., & Strickland, D.K. 1997, *MNRAS*, 286, 626
 Read, A.M., & Pietsch, W. 2001, *A&A*, 373, 473
 Sarazin, C.L., Irwin, J.A., & Bregman, J.N. 2001, *ApJ*, 556, 533
 Scargle, D. 1982, *ApJ*, 263, 835
 Soria, R., & Kong, A.K.H. 2003, *A&A*, submitted
 Soria, R., & Wu, K. 2003, *A&A*, submitted
 Strickland, D.K., Colbert, E.J.M., Heckman, T.M., Weaver, K.A., Dahlem, M., & Stevens, I.R. 2001, *ApJ*, 560, 707
 Swartz, D.A., Ghosh, K.K., Suleimanov, V., Tennant, A.F., Wu, K. 2002, *ApJ*, 574, 382
 Tully, R. 1988, *Nearby Galaxies Catalog* (Cambridge: Cambridge Univ. Press)
 van Teeseling, A., & King, A.R. 1998, *A&A*, 338, 957

TABLE 1
BEST-FITTING SPECTRAL PARAMETERS

Date	N_H ($\times 10^{21} \text{ cm}^{-2}$)	kT (eV)	L_X ^a	L_{bol} ^b	$\chi^2_{\nu}/\text{d.o.f}$
1992 May 26	$1.08^{+0.31}_{-0.31}$	$48.7^{+9.1}_{-9.1}$	10	20	0.74/11
2000 Dec 26	$1.50^{+0.32}_{-0.20}$	$56.8^{+2.2}_{-3.3}$	9	15	1.07/39
2001 Jan 1	$0.53^{+0.20}_{-0.07}$	$66.5^{+2.4}_{-3.6}$	1.1	1.9	1.15/48

Note. — All quoted uncertainties are 1σ .

^a0.2–2 keV luminosity ($\times 10^{38} \text{ erg s}^{-1}$), assuming a distance of 2 Mpc

^bBolometric luminosity ($\times 10^{38} \text{ erg s}^{-1}$)




FIG. 1.— Energy spectra of XMMU J005510.7-373855 (see Table 1 for spectral parameters). From left to right: *ROSAT* PSPC spectrum on 1992 May/June (high state), *XMM-Newton* spectrum on 2000 December 26 (high state) and *XMM-Newton* spectrum on 2001 January 1 (low state). For the *XMM-Newton* spectra, pn, MOS1, and MOS2 data are marked as triangles, solid squares and circles.




FIG. 2.— Long-term lightcurve of the supersoft ULX from 1991 to 2001 (circles: *ROSAT* PSPC; squares: *ROSAT* HRI; triangles: *XMM-Newton*). The 0.2–2 keV luminosity is from spectral fits in Table 1, while the 3σ upper limits are assuming a blackbody model with $N_H = 10^{21} \text{ cm}^{-2}$ and $kT = 58 \text{ eV}$.




FIG. 3.— a) Lomb-Scargle periodogram of XMMU J005510.7-373855 as obtained by *XMM-Newton* on 2001 January. The horizontal dotted line is the 99.9% confidence level; b) Folded lightcurve of the “low” state (2001 January) data on a period of 5.4 hr; c) Folded lightcurve of the “high” state (2000 December) data on a period of 5.4 hr. T_0 of both lightcurves is set at the time of the first data point.